方知库
Eco-Environmental
Knowledge Web

环境科学

ENVIRONMENTAL SCIENCE

ISSN 0250-3301 CODEN HCKHDV
HUANJING KEXUE

- 主办 中国科学院生态环境研究中心
- 出版 科学出版社

2020

Vol.41 No.11
第41卷 第11期

目次

中国工业源挥发性有机物排放清单 梁小明, 孙西勃, 徐建铁, 叶代启, 陈来国(4767)

中国水泥排放清单及分布特征 汤铃, 薛晓达, 伯鑫, 郭静, 王鹏, 翟文慧, 崔维庚, 汪寿阳, 李时蓓, 董广霞(4776)

环境空气 PM_{2.5} 化学组分监测数据审核指标的建立:以长三角地区为例 张哲, 乔利平, 周敏, 朱书慧, 郭会琴, 王红丽, 楼晟荣, 陶士康, 陈长虹(4786)

南京江北新区大气颗粒物化学组分的粒径分布特征 张毓秀, 于兴娜, 刘偲嘉, 安俊琳, 张程(4803)

郑州市 PM_{2.5} 化学组分的季节变化特征及来源解析 张剑飞, 姜楠, 段时光, 孙有昌, 郝祺, 张瑞芹(4813)

山西大学城 PM_{2.5} 中元素特征、来源及健康风险评估 李雪梅, 牟玲, 田妹, 郑利荣, 李杨勇(4825)

2000~2019年中国 PM_{2.5} 时空演化特征 夏晓圣, 汪军红, 宋伟东, 程先富(4832)

北京地区偏南风 and 偏东风条件下污染特征差异 尹晓梅, 乔林, 朱晓婉, 郭恒, 刘湘雪, 熊亚军(4844)

基于天气背景天津大气污染输送特征分析 蔡子颖, 杨旭, 韩素芹, 姚青, 刘敬乐(4855)

基于天气分型的上甸子大气本底站臭氧污染气象条件 吴进, 李琛, 马志强, 孙兆彬, 朱晓婉, 董璠(4864)

基于环境空气中 VOCs 在线监测数据精准识别化工园区 VOCs 排放源 孙佳宁, 谢绍东(4874)

芜湖市大气挥发性有机物污染特征、大气反应活性及源解析 高亢, 章慧, 刘梦迪, 赵晓宇, 吴婷, 贾其娜(4885)

密云水库上游白河地表水质对不同空间尺度景观格局特征的响应 张微微, 李晓娜, 王超, 赵春桥, 史瑞双(4895)

北京密云水库流域水体夏季 DOC 和 DIC 质量浓度及同位素组成初探 陈吉吉, 郭婧, 徐苏士, 陶蕾, 荆红卫(4905)

2009~2018年太湖湖泛强度变化及其影响因素 辛华荣, 朱广伟, 王雷松, 李骏, 陆隽(4914)

雷州半岛岭北地区地下水水文地球化学特征 张宏鑫, 吴亚, 罗炜宇, 陈雯, 刘怀庆(4924)

基于多环境介质氮素和同位素的滦平盆地地下水硝酸盐来源示踪 孙厚云, 卫晓锋, 贾凤超, 李多杰, 李健, 李霞, 殷志强(4936)

会仙岩溶湿地水体金属元素分布与健康风险评价 李军, 邹胜章, 梁永平, 林永生, 周长松, 赵一(4948)

太白山自然保护区水体 CDOM 吸收与三维荧光特征 张文浩, 赵铎霖, 王晓毓, 郝强州, 李喜梅, 高鹏程(4958)

强降雨对平原河网区入湖河道氮、磷影响 连心桥, 朱广伟, 杨文斌, 朱梦圆, 许海(4970)

长江流域淡水生态系统内分泌干扰物、药物和个人护理品的风险排序 苏超, 崔严(4981)

嘉兴市城市河网区多环芳烃污染源解析及生态风险评价 杨延梅, 赵航晨, 孟睿, 何连生, 赵昊, 杜士林, 丁婷婷(4989)

污水处理厂对漓江桂林市河段淡水生物中微塑料累积的影响 谢元沅, 张清科, 张漓杉, 钟山, 刘俊勇, 侯晓映, 王华, 冯静培, 梁祖鸿, 陈舞阳(4999)

村镇生活污水处理设施抗生素浓度分布规律 卢亚楠, 郭雅妮, 王坤, 陆伟东, 王拓, 柴玉峰, 柳蒙蒙, 钟慧, 陈梅雪, 魏源送(5008)

冬季小流域水体微生物多样性及影响因素 朱金山, 秦海兰, 孙启耀, 王宝珍, 高润霞, 郭锐利, 李武斌(5016)

水库热分层期沉积物中有机物特性及对好氧反硝化细菌脱氮的影响 王彤, 文刚, 黄廷林, 司凡, 王晨旭, 路林超, 林薇(5027)

长三角地区 MBBR 泥膜复合污水厂低温季节微生物多样性分析 韩文杰, 吴迪, 周家中, 何强, 阚渝姣(5037)

类群划分方法对南亚热带水库夏季浮游植物群落与环境响应关系的影响 郭芳, 顾继光, 赵剑, 余华章, 肖利娟(5050)

基于决策树的城市黑臭水体遥感分级 李玲玲, 李云梅, 吕恒, 徐杰, 杨子谦, 毕顺, 许佳峰(5060)

纳米零价铁强化微生物电催化-厌氧膜生物组合反应器抗膜污染能力及其调控机制 潘阳, 牛承鑫, 支忠祥, 王建辉, 陆雪琴, 戴金山, 甄广印, 孙彤彤, 康娜英(5073)

温度和基质浓度对厌氧氨氧化工艺中 N₂O 释放的影响 王丝可, 于恒, 左剑恶(5082)

以污泥发酵液与尾水混合液为补给碳源的 A²/O 污染物去除特性 荣懿, 邵劲博, 于优城, 石恒, 金鹏康(5089)

不同曝气比对连续流亚硝化颗粒污泥运行的影响 张杰, 劳会妹, 李冬, 魏子清(5097)

中温厌氧消化与高温厌氧消化对污泥重金属风险及稳定性的影响 田中科, 王芬, 闫钊(5106)

基于 PMF 模型及地统计的土壤重金属健康风险定量评价 韩琳, 徐夕博(5114)

再开发利用工业场地土壤重金属含量分布及生态风险 沈城, 刘霞雯, 吴健, 黄沈发, 王敏, 黄波涛(5125)

有机改性蒙脱土对苯的吸附机制和影响因子 侯静雯, 施维林, 高娟(5133)

硫酸亚铁和硝酸铁施用对根际土壤-水稻系统中镉迁移的影响 李义纯, 陈勇, 唐明灯, 李林峰, 林晓扬, 王艳红, 许迪豪, 艾绍英(5143)

叶用油菜和孔雀草间作对植物生长和镉累积的影响 闫秀秀, 徐应明, 王林, 陶雪莹, 孙约兵, 梁学峰(5151)

重金属固定细菌调控土壤铁氧化物阻控小麦 Cd 吸收效应 王晓宇, 韩辉(5160)

油-稻轮作模式下修复材料对土壤铜修复的持续性影响 赵颖颖, 李虹川, 胡宏祥, 薛中俊, 刘娜, 吴孟君(5168)

氮肥使用对北方夏玉米季氨挥发的影响 张薇, 倪邦, 许秀春, 杨轩, 孟凡乔(5176)

生物炭影响下土壤呼吸温度敏感性及其细菌群落结构的变化 曹坤坤, 张沙沙, 胡学玉, 黄洋(5185)

铜尾矿白羊草重金属含量对叶际和根际真菌群落的影响 贾彤, 郭婷艳, 王瑞宏, 柴宝峰(5193)

典型水环境微生物源异味物研究进展 张海涵, 苗雨甜, 黄廷林, 刘凯文, 刘祥, 黄鑫, 杨尚业, 司凡, 李程遥(5201)

广告:国家微生物科学数据中心 (前插)

《环境科学》征订启事(4785) 《环境科学》征稿简则(4947) 信息(5015, 5124, 5132)

长江流域淡水生态系统内分泌干扰物、药物和个人护理品的风险排序

苏超¹, 崔严²

(1. 山西大学黄土高原研究所, 太原 030006; 2. 山西大学环境与资源学院, 太原 030006)

摘要: 作为新兴污染物, 内分泌干扰物、药物和个人护理品因其排放对水环境安全产生的风险而引起了广泛关注. 本文以长江流域为例, 以 69 种内分泌干扰物、药物和个人护理品的环境暴露数据和生态毒性数据为基础, 计算它们的生态风险并进行高低排序, 以识别最优先控制的污染物. 结果表明, 内分泌干扰物和个人护理品对水生生态系统的风险较高, 如雌酮 (estrone)、雌三醇 (estriol)、17 β -雌二醇 (17 β -estradiol)、双酚 S (bisphenol S)、阿特拉津 (atrazine)、三氯卡班 (triclocarban) 和三氯生 (triclosan); 而药物的风险低 500 倍或更多. 相较而言, 药物中风险较高的属抗生素类. 湘江和洪湖及其周边河流内多种污染物的浓度呈现出最高值, 是长江流域污染最重的区域, 其次是太湖、洞庭湖和长江三角洲地区. 与藻类和虫类相比, 鱼类是对内分泌干扰物、药物和个人护理品最敏感的物种, 尤其对 17 α -乙炔雌二醇 (17 α -ethynylestradiol) 和 17 β -雌二醇. 本文的研究结果可对我国化学品的监测和管理提供数据支撑.

关键词: 长江流域; 内分泌干扰物 (EDCs); 药物; 个人护理品; 风险排序

中图分类号: X52; X820.4 文献标识码: A 文章编号: 0250-3301(2020)11-4981-08 DOI: 10.13227/j.hjxx.202003210

Risk Ranking of Endocrine Disrupting Compounds, Pharmaceuticals, and Personal Care Products in the Aquatic Environment of the Yangtze River Basin

SU Chao¹, CUI Yan²

(1. Institute of Loess Plateau, Shanxi University, Taiyuan 030006, China; 2. College of Environmental & Resource Sciences, Shanxi University, Taiyuan 030006, China)

Abstract: Endocrine disrupting compounds (EDCs), pharmaceuticals, and personal care products (PPCPs) have been of great concern as emerging contaminants of aquatic environments. In this study, the risks of EDCs and PPCPs in the Yangtze River Basin were ranked, based on their environmental exposure datasets and ecotoxicity datasets. The prioritized chemicals were then identified. The results found that EDCs and personal care products were deemed to represent higher risks to aquatic organisms, for example, estrone, estriol, 17 β -estradiol, bisphenol S, atrazine, triclocarban, and triclosan, while the risks of pharmaceuticals were 500-fold lower. Among the pharmaceuticals, antibiotics posed relatively higher risks. The Xiangjiang River and Honghu Lake with its surrounding rivers were hotspots where more chemicals were found with the highest concentrations, followed by Taihu Lake, Dongting Lake and the Yangtze River Delta. When comparing with algae and worms, fish were the most sensitive taxa to 17 β -estradiol and 17 α -ethynylestradiol. The results of this work provide sound guidance for the future monitoring and management of chemicals in China.

Key words: Yangtze River Basin; endocrine disrupting compounds (EDCs); pharmaceuticals; personal care products; risk ranking

近些年来, 内分泌干扰物 (endocrine disrupting compounds, EDCs)、药物和个人护理品 (pharmaceuticals and personal care products, PPCPs) 作为新兴污染物受到了众多环保工作者的广泛关注^[1-8]. 药物主要用于预防或治疗人畜疾病, 包括抗生素、止痛药和血脂调节剂等, 其中抗生素因其广泛应用而受到特别关注^[4,7,8]. 在治疗期间和治疗之后, 动物和人类排泄代谢部分或完整的药物, 而这些药物最终直接或间接地排放到环境中^[8]. 个人护理品 (如消毒剂、香水、洗发水和防晒霜等) 常被用于改善人们的生活质量^[7]. 内分泌干扰物是一类即使在浓度极低的情况下, 依然会影响动物和人类的生殖、发育过程或干扰其内分泌系统的化学物质^[5,9]; 它们通常来自于人畜的排泄物 (如合成类固醇激素的药物和合成雄激素) 或人造产品 (如塑料制品)^[10].

目前 EDCs 和 PPCPs 在世界范围内被广泛使用, 并且极易排放到环境中. 这些物质大多数易溶于水, 因此 EDCs 和 PPCPs 在世界各地的淡水环境中被普遍检出^[1,11-13]. 而且, 这些物质中的大多数具有持久性有机污染物的特点, 即持久性、毒性和生物富集性. 因此, EDCs 和 PPCPs 对生态系统健康的不良影响引起了广泛关注.

由于我国的快速发展和人口激增, 我国成为 EDCs 和 PPCPs 的最大生产国和消费国. 据估算, 2011 年我国的药品消费量约为 200 万 t, 其中抗生

收稿日期: 2020-03-21; 修订日期: 2020-05-09

基金项目: 山西省高等学校科技创新项目 (2019L0017); 山西省高等学校“服务产业创新学科群建设计划”项目 (20181401); 国家自然科学基金项目 (41771548)

作者简介: 苏超 (1989 ~), 女, 博士研究生, 讲师, 主要研究方向为污染物的风险评估及环境行为模拟, E-mail: suchao@sxu.edu.cn

素的消费量超过了 18 万 t,人均消费量远远超过美国^[14,15],这间接导致了抗生素在水环境中的普遍检出.另外,其他种类的药物、个人护理品和内分泌干扰物也在水环境中普遍存在.鉴于众多种类的化学物质被检出,那么,分析每种化学物质的相对风险,并确定需要优先控制的污染物已显得非常必要.长江流域是我国的第一大流域,流域面积约占中国陆地总面积的 1/5,水资源总量超过全国河流径流量的 1/3,人口和经济总量均超过全国的 40%^[16,17].因此,本文以长江流域为研究区,以文献搜集到的 EDCs 和 PPCPs 的环境暴露浓度和生态毒性浓度为数据基础,计算每种污染物的生态风险并确定优先控制的污染物,分析长江流域的热点污染区域,识别最敏感的生物群,以期为化学品的风险管理提供数据基础.

1 材料与方 法

1.1 污染物筛选及环境暴露数据收集

污染物的选择由其可获得性和环境暴露数据的质量决定.中文文献来自中国知网、维普和万方数据库,同时包括硕士毕业论文和博士毕业论文.英文文献来自于 Web of Science 和 Google Scholar 两个数据库.首先,以“内分泌干扰物 & 水/河/湖 & 中国”、“药物 & 水/河/湖 & 中国”、“抗生素 & 水/河/湖 & 中国”和“个人护理品 & 水/河/湖 & 中国”为主题关键词进行文献搜索.之后筛选出属于英文 SCI 期刊和中文核心期刊(包括所有硕士和博士毕业论文)且采样位置位于长江流域的文献进行统计.本文只统计采样时间在 2014 年 1 月至 2019 年 9 月的文献.若文献里的浓度数据模糊、采样时间不符、地理信息不匹配或不清等,则予以删除.对于检索到的数据,若文献里明确说明数据有异常值,则删除异常值,否则保留全部数据.对于任何一种污染物而言,纳入研究的物质必须至少包含 3 个采样点的数据.此外,本文不考虑排污河和水库的浓度数据.当污染物的暴露浓度低于检出限(LOD)时,其浓度用 $LOD/\sqrt{2}$ 表示^[14,18,19].

1.2 生态毒理数据收集和筛选

污染物的生态毒理数据主要在美国环保署(USEPA)网站的 ECOTOX 数据库里下载得到,同时包括了从文献中检索到的毒性数据.这里的毒性数据库包括了大多数物种(如藻类、鱼类、植物和软体动物等)的效应浓度.首先,从数据库里中提取出环境介质为淡水的毒性数据.其次,本文同时考虑了毒性数据的致死和亚致死效应浓度.对任一物种, EC_{50} (effect concentration 50) 和 LC_{50} (lethal concentration

50) 是最优先选择的评价终点,其次为 EC_{xx} 和 LC_{xx} (如 EC_{20} 、 EC_{10} 和 LC_{20} 等),最后为最低效应浓度(LOEC)^[14,18,19].对任一物种,若同一评价终点下有多个值,那么选取最低值.随后,只有中国的本土物种和标准测试物种数据被保留.最后,对于任何一种污染物而言,纳入研究的物质必须至少包含 3 条毒性数据.

1.3 风险评价

收集完成污染物的毒性数据集和暴露数据集后,它们反映出的信息便可以分别绘制出来.对每一种污染物的每一个数据集,需要计算出它们的中值.那么,毒性数据集和暴露数据集的中值的接近程度可以用一个比例来表示,如公式(1)所示,这个比例被定义为污染物的风险 R ^[20-23].那么,比值越大,表示污染物的风险越大.

$$R = \frac{M_w}{M_r} \quad (1)$$

式中, M_w 表示暴露浓度的中值 ($ng \cdot L^{-1}$), M_r 表示毒性数据的中值 ($ng \cdot L^{-1}$).

此外,这个方法还可用来比较特定污染物对不同生物群体(如藻类和鱼类等)的相对风险,从而识别最敏感的生物群体.

2 数据概况与范围

根据上述数据检索规则,本文共综述了 257 篇文献,其中 204 篇由于浓度数据模糊、采样时间不符、地理信息不匹配或不清等原因不予考虑.最后进行统计的文献中,包括 19 篇中文和 34 篇英文文献.数据统计完成后发现,有 117 种 EDCs 和 PPCPs 至少被报道了一次;之后,由于缺乏充分的暴露数据或毒性数据,共有 69 种 EDCs 和 PPCPs 参与风险排序,包括 19 种内分泌干扰物,41 种药物和 9 种个人护理品.初步分析发现,大多数的药物为抗生素,包括磺胺类、四环素类、大环内酯类和氟喹诺酮类.个人护理品中,作为广谱杀菌剂的三氯生(TCS)和三氯卡班(TCC)是报道最多的两种.此外,酚类 EDCs(如双酚 A 和双酚 S)和甾体类 EDCs(如 17β -雌二醇和 17α -乙炔雌二醇)是最主要的两类内分泌干扰物.

3 结果与讨论

3.1 淡水环境中污染物的风险排序

基于污染物对所有生物种的风险,本文得到了 69 种污染物的相对风险排序结果(图 1).从图 1 中可看出,整体上,内分泌干扰物(橘色)和个人护理品(蓝色)表现出较高风险,而药物(绿色)呈现出较低风险.风险最高的 10 种物质依次是雌酮(E1)、雌

三醇(E3)、17 β -雌二醇(β E2)、双酚 S(BPS)、阿特拉津(ATZ)、17 α -乙炔雌二醇(EE2)、对-特辛基苯酚(4-t-OP)、三氯卡班(TCC)、三氯生(TCS)和双酚 AF(BPAF),即以内分泌干扰物和个人护理品为主. 相比之下,苯甲酮-2(BP-2,排名 59)和对-羟基苯甲酸乙酯(EtP,排名 61)对生物体的风险较低. 与雌酮的风险相比,药物的风险低 500 倍或更多(图 2). 药物中,风险较高的是抗生素类,如洛美沙星(LOM)、磺胺甲噁唑(SMX)、红霉素(ETM)、强力霉素(DOX)、四环素(TC)和罗红霉素(RTM)等,以及氟西汀(FLX); 然而,氯霉素(CHP)、吉非贝齐(GFB)、美托洛尔(MTP)、吡哌美辛(IMC)、磺胺噻

唑(STZ)、萘普生(NAP)、对-乙酰氨基酚(ACE)和苯扎贝特(BZB)等污染物处于较低风险. Li 等^[24]的评价结果认为,在众多药物中,双氯芬酸(DIC)、红霉素、青霉素、阿奇霉素和阿莫西林等抗生素类药物是中国淡水环境风险最高的物质; 与之相比,本文发现,在长江流域,上述内分泌干扰物和个人护理品的风险总体上高于双氯芬酸、红霉素和阿莫西林等药物的风险. 此外,观察图 1 中毒性数据集的较小值和暴露数据集的较大值,发现多种污染物的暴露数据和毒性数据表现出较大比例的重叠分布,如雌酮、17 β -雌二醇、双酚 S、阿特拉津、17 α -乙炔雌二醇和对-特辛基苯酚等,表明多种水生生物处于危险状态.

1. estrone (E1), 2. estriol (E3), 3. 17 β -estradiol (β E2), 4. bisphenol S (BPS), 5. atrazine (ATZ), 6. 17 α -ethynylestradiol (EE2), 7. 4-tert-octylphenol (4-t-OP), 8. triclocarban (TCC), 9. triclosan (TCS), 10. bisphenol AF (BPAF), 11. benzophenone-4 (BP-4), 12. benzophenone (BP), 13. bisphenol A (BPA), 14. lomefloxacin (LOM), 15. 4-nonylphenol (4-NP), 16. testosterone (TES), 17. sulfamethoxazole (SMX), 18. 4-tert-butylphenol (4-TBP), 19. erythromycin (ETM), 20. 4-octylphenol (4-OP), 21. doxycycline (DOX), 22. fluoxetine (FLX), 23. tetracycline (TC), 24. phantolide (PHE), 25. roxithromycin (RTM), 26. ibuprofen (IBU), 27. caffeine (CAF), 28. sulfadiazine (SDZ), 29. benzophenone-3 (BP-3), 30. diazepam (DZM), 31. oxytetracycline (OTC), 32. clofibrate (CA), 33. lincomycin (LCM), 34. diethylstilbestrol (DES), 35. p-hydroxybenzoic acid (PHBA), 36. butyl paraben (BuP), 37. sulfamethazine (SMZ), 38. ofloxacin (OFL), 39. norfloxacin (NOR), 40. mefenamic acid (MA), 41. 4-butylphenol (4-BP), 42. diphenhydramine (DHM), 43. phenazone (PHO), 44. carbamazepine (CBZ), 45. diethyltoluamide (DEET), 46. climbazole (CLM), 47. sulfadiazine (SDZ), 48. diclofenac (DIC), 49. ciprofloxacin (CIP), 50. florfenicol (FFC), 51. methyl paraben (MeP), 52. propyl paraben (PrP), 53. amoxicillin (AMX), 54. chlortetracycline (CTC), 55. enrofloxacin (ENR), 56. thiamphenicol (TAP), 57. trimethoprim (TMP), 58. tylosin tartrate (TS), 59. benzophenone-2 (BP-2), 60. paracetamol (PRC), 61. ethyl paraben (EtP), 62. bezafibrate (BZB), 63. acetaminophen (ACE), 64. naproxen (NAP), 65. sulfathiazole (STZ), 66. indomethacin (IMC), 67. metoprolol (MTP), 68. gemfibrozil (GFB), 69. chloramphenicol (CHP); 每种污染物的左边列和圆形图例表示毒性数据,右边列和菱形图例表示环境暴露数据;每一个数据集的中值以空心圆表示

图 1 基于配对的毒性数据和环境暴露数据的 69 种污染物的风险排序

Fig. 1 Risk ranking of 69 chemicals using paired datasets of all the collected ecotoxicity data and measured freshwater concentrations

雌酮对长江流域的生物体表现出最大风险,风险值 R 为 0.040,其次为雌三醇($R=0.034$)和 17 β -雌二醇($R=0.0079$),见图 2. 长江流域的河(湖)水中,雌酮的暴露浓度的中值为 $3.00 \text{ ng}\cdot\text{L}^{-1}$. 雌酮的最低效应浓度(也即最敏感)为 $3.30 \text{ ng}\cdot\text{L}^{-1}$,为幼年虹鳟鱼(*Oncorhynchus mykiss*)的 LOEC 值^[25];对斑马鱼(*Danio rerio*)而言,暴露 18 d 的 EC_{50} 值低至 $78.00 \text{ ng}\cdot\text{L}^{-1}$ ^[26]. 雌酮和雌三醇是常见且流行的类固醇化合物,被广泛用为口服避孕药和人类、牲畜疾病的治疗药物,因此常作为人和牲畜的排泄物进入

环境^[27]. 正是由于它们的广泛使用,它们在世界各地的地表水中的检出率超出 70%,浓度范围从 $\text{ng}\cdot\text{L}^{-1}$ 至 $\mu\text{g}\cdot\text{L}^{-1}$ 不等^[5]. 长江流域作为我国第一大流域,人口总量超过全国的 40%^[16,17],养殖业发达,导致雌酮和雌三醇的暴露浓度范围分别为 $\text{ND}\sim 84.00 \text{ ng}\cdot\text{L}^{-1}$ 和 $\text{ND}\sim 82.00 \text{ ng}\cdot\text{L}^{-1}$. 此外,内分泌干扰物和个人护理品被广泛用于人类的各种消费品中,随着长江流域城市化和现代化水平的快速发展,产生了大量含有内分泌干扰物和个人护理品的生活污水、工业废水和养殖废水,而这些物质在废水处理

图2 基于污染物暴露数据和毒性数据中值的风险值排序结果

Fig. 2 Risk ranking of the chemicals by plotting the ratios of the median environmental concentrations and median effect concentrations

过程中并不能被完全去除,因此这些物质可能经污水处理厂排放、养殖废水排放和地表径流等方式进入长江水域,从而造成内分泌干扰物和个人护理品

在长江流域的高暴露水平^[28].加之它们的效应浓度较低,因此内分泌干扰物和个人护理品呈现出较高风险.

表1 EDCs 和 PPCPs 的最高浓度分析¹⁾/ng·L⁻¹

Table 1 Rivers/lakes with the highest reported concentrations of EDCs and PPCPs/ng·L⁻¹

排名	物质	最高浓度	相应位置	排名	物质	最高浓度	相应位置
1	E1	83.52	洪湖 ^[32]	36	BuP	9.50	湘江 ^[49]
2	E3	81.60	三峡库区 ^[40]	37	SMZ	715.30	洪湖周边河流 ^[45]
3	βE2	58.94	洪湖 ^[32]	38	OFL	203.70	洪湖周边河流 ^[45]
4	BPS	1 569.00	太湖 ^[33]	39	NOR	142.00	东荆河 ^[41]
5	ATZ	496.00	京杭运河 ^[34]	40	MA	324.03	长江三角洲 ^[46]
6	EE2	43.93	洪湖 ^[32]	41	4-BP	282.40	太湖 ^[42]
7	4-t-OP	156.10	湘江 ^[9]	42	DHM	0.89	汉江 ^[35]
8	TCC	12.66	汉江 ^[35]	43	PHO	22.50	东洞庭湖 ^[44]
9	TCS	65.60	长江中游 ^[36]	44	CBZ	145.95	长江三角洲 ^[46]
10	BPAF	140.00	太湖 ^[37]	45	DEET	574.75	长江三角洲 ^[46]
11	BP-4	131.00	黄浦江 ^[38]	46	CLM	100.00	湘江 ^[47]
12	BP	712.59	苏州河 ^[38]	47	SDZ	499.50	洪湖周边河流 ^[45]
13	BPA	771.50	湘江 ^[9]	48	DIC	230.05	东洞庭湖 ^[44]
14	LOM	14.60	太湖 ^[39]	49	CIP	106.20	洪湖周边河流 ^[45]
15	4-NP	858.00	长江(南京段) ^[40]	50	FFC	63.50	太湖 ^[39]
16	TES	70.40	湘江 ^[9]	51	MeP	3 173.90	湘江 ^[49]
17	SMX	173.46	三峡库区 ^[41]	52	PtP	1 040.40	湘江 ^[49]
18	4-TBP	1 121.00	长江(南京段) ^[40]	53	AMX	710.00	湘江 ^[47]
19	ETM	982.00	东荆河 ^[41]	54	CTC	876.20	洪湖周边河流 ^[45]
20	4-OP	305.10	太湖 ^[42]	55	ENR	96.55	鄱阳湖 ^[43]
21	DOX	23.93	鄱阳湖 ^[43]	56	TAP	12.00	湘江 ^[47]
22	FLX	40.20	东洞庭湖 ^[44]	57	TMP	209.69	三峡库区 ^[50]
23	TC	1454.80	洪湖周边河流 ^[45]	58	TS	19.50	三峡库区 ^[51]
24	PHE	66.41	长江三角洲 ^[46]	59	BP-2	34.70	黄浦江 ^[38]
25	RTM	190.00	湘江 ^[47]	60	PRC	65.50	汉江 ^[35]
26	IBU	320.00	湘江 ^[47]	61	EtP	87.00	湘江 ^[49]
27	CAF	174.40	东洞庭湖 ^[44]	62	BZB	0.05	洞庭湖 ^[52]
28	SMD	172.40	洪湖 ^[45]	63	ACE	3.88	洞庭湖 ^[52]
29	BP-3	5 013.07	黄浦江 ^[38]	64	NAP	125.71	长江三角洲 ^[46]
30	DZM	104.92	长江三角洲 ^[46]	65	STZ	8.18	太湖 ^[39]
31	OTC	2 796.60	洪湖周边河流 ^[45]	66	IMC	2.70	湘江 ^[47]
32	CA	13.00	太湖 ^[48]	67	MTP	1.60	东洞庭湖 ^[44]
33	LCM	53.80	太湖 ^[39]	68	GFB	30.50	洞庭湖 ^[52]
34	DES	9.60	湘江 ^[9]	69	CHP	6.10	湘江 ^[47]
35	PHBA	2 380.00	长江中游 ^[36]				

1) 表中污染物的排名和图1的一致

在风险较高的药物中,氟西汀为临床广泛应用的抗抑郁药物,其环境浓度范围为 $\text{ND} \sim 40.20 \text{ ng}\cdot\text{L}^{-1}$,中值为 $1.60 \text{ ng}\cdot\text{L}^{-1}$;其效应浓度的中值为 $169.81 \mu\text{g}\cdot\text{L}^{-1}$.洛美沙星、磺胺甲噁唑、红霉素、强力霉素、四环素和罗红霉素都是常用的抗生素.在长江流域,洛美沙星、磺胺甲噁唑、红霉素、强力霉素、四环素和罗红霉素的环境浓度范围分别为 $0.50 \sim 14.60$ 、 $\text{ND} \sim 173.46$ 、 $\text{ND} \sim 982.00$ 、 $\text{ND} \sim 23.93$ 、 $\text{ND} \sim 1454.80$ 和 $\text{ND} \sim 190.00 \text{ ng}\cdot\text{L}^{-1}$,中值分别为 2.85 、 11.80 、 5.80 、 2.20 、 8.89 和 $8.10 \text{ ng}\cdot\text{L}^{-1}$.它们效应浓度的中值分别为 141.50 、 775.00 、 430.00 、 215.50 、 1000.00 和 $1000.00 \text{ ng}\cdot\text{L}^{-1}$.文献报道的洛美沙星的效应浓度的最低值为 $30.00 \mu\text{g}\cdot\text{L}^{-1}$,为暴露 7 d 的浮萍 (*Lemna gibba*) 的 LOEC 值,该浓度会影响浮萍的繁殖^[29].磺胺甲噁唑的效应浓度的最低值为 $3.36 \mu\text{g}\cdot\text{L}^{-1}$,为暴露 7 d 的浮萍的 EC_{50} 值.事实上,磺胺类、氟喹诺酮类和大环内酯类的抗生素在中国的消费量约占抗生素总消费量的 12%、15% 和 20%^[14,30,31].然而,现有的研究发现抗生素对水生生物体是相对安全的,而且本文没有考虑生物体对抗生素的抗药性.

3.2 热点污染区域分析

每种污染物的最高浓度及其相应位置见表 1.

从中可知,湘江是长江流域污染最重的河流,其中 15 种污染物的最高浓度均位于此,包括对-特辛基苯酚^[9]、双酚 A (BPA)^[9]、睾酮 (TES)^[9]、罗红霉素^[47]、布洛芬 (IBU)^[47]、己烯雌酚 (DES)^[49]、对-羟基苯甲酸丁酯 (BuP)^[49]、氯咪巴唑 (CLM)^[47]、对-羟基苯甲酸甲酯 (MeP)^[49]、对-羟基苯甲酸丙酯 (PrP)^[49]、阿莫西林 (AMX)^[47]、甲砒霉素 (TAP)^[47]、对-羟基苯甲酸乙酯 (EtP)^[49]、吡喹酮^[47] 和氯霉素^[47].污染热点区域其次是洪湖及其周边河流^[32,45]、太湖^[33,37,39,42,48]、洞庭湖^[44,52] 和长江三角洲地区^[46].另外,从污染物的风险高低顺序可看出,洪湖及其周边河流、太湖和湘江的多种污染物处于较高风险(表 1).

湘江是长江流域的一个主要支流,长约 856 km,流域人口大于 4 千万,被誉为湖南省的“母亲河”,是湖南省的重要经济纽带^[9,47].湘江流域包含着许多养殖区、工业区、耕作区、医院和药品生产企业^[47].近些年来,随着人口增长、经济发展和快速的城市化过程,湘江流域的水污染现象日益突出.有研究表明,工业加工助剂是对-特辛基苯酚和双酚 A 的主要来源^[9];此外,污水处理厂污水的低处理效率和生活污水的直接排放也是上述污染物的主要来源^[47,49].

图 3 部分内分泌干扰物和药物对藻类、鱼类和虫类的风险排序

Fig. 3 Risk ratio ranking of some EDCs and pharmaceuticals by algae, fish, and worms

洪湖是长江中游的一个典型浅水湖,水产养殖业是其支柱产业,围网养鱼现象在洪湖周边普遍存在;而围栏和围网是围网养殖业的主要方式,因此成为内分泌干扰物的主要来源^[32].另外,洪湖周边的人口密集,畜牧业也非常发达(>8 百万头猪),有大量生活污水和畜牧业污水排入湖内,而内分泌干扰物和抗生素被广泛用于人类生活和水产养殖业、畜牧业中,因此洪湖及其周边河流内的多种内分泌干扰物和抗生素呈现出最高水平^[45].

3.3 敏感物种识别

根据图 1 的风险排序结果,本文选择排名前 40 的污染物,并筛选出那些毒性数据包含共同生物群的污染物,来进行敏感物种分析.这些污染物是阿特拉津、红霉素、磺胺甲噁唑、对-壬基苯酚(4-NP)、双酚 A、氟西汀、四环素、17 β -雌二醇和 17 α -乙炔雌二醇,之后分别计算出这些污染物对藻类、鱼类和虫类的风险,计算得到的风险排序结果见图 3.从中可知,内分泌干扰物对藻类、鱼类和虫类的风险都较高.具体地,阿特拉津对藻类和虫类的风险较高,风险值几乎处于同一量级,但是对鱼类的风险较低(图 3).17 α -乙炔雌二醇和 17 β -雌二醇(分别排名 1 和 2)对鱼类的风险远远高于其他污染物,但是对藻类(分别排名 8 和 9)和虫类(分别排名 5 和 6)的风险极低.另一区别是,药物(红霉素和磺胺甲噁唑)对藻类和虫类呈现出较高风险,而对鱼类的风险很低.

从风险值来看,与藻类和虫类相比,鱼类是最敏感的生物,尤其是对 17 α -乙炔雌二醇和 17 β -雌二醇,风险值分别为 0.27 和 0.10[图 3(b)],约为藻类和虫类的 100 000 倍.与这两种污染物相比,其他污染物对鱼类的风险低 1 000 倍或更多.藻类是对阿特拉津(一种除草剂)最敏感的物种,而鱼类是最不敏感的.相比之下,因为较高的效应浓度,虫类是最不敏感的物种.

4 结论

(1) 内分泌干扰物和个人护理品的生态风险相对较高,而药物的风险较低;药物中,风险较高的属抗生素类.

(2) 湘江流域是长江流域污染最重的区域,其次是洪湖及其周边河流、太湖、洞庭湖和长江三角洲地区.

(3) 与藻类和虫类相比,鱼类是最敏感的物种,尤其对 17 α -乙炔雌二醇和 17 β -雌二醇,风险值约为藻类和虫类的 100 000 倍.相比之下,虫类是最不敏感的物种.

(4) 长江流域是我国的第一大流域,对其淡水生态系统的新兴污染物开展风险评估及排序,可为化学品的风险管理提供指导.

参考文献:

- [1] García-Galán M J, Arashiro L, Santos L H M L M, *et al.* Fate of priority pharmaceuticals and their main metabolites and transformation products in microalgae-based wastewater treatment systems [J]. *Journal of Hazardous Materials*, 2020, **390**: 121771, doi: 10.1016/j.jhazmat.2019.121771.
- [2] La Merrill M A, Vandenberg L N, Smith M T, *et al.* Consensus on the key characteristics of endocrine-disrupting chemicals as a basis for hazard identification [J]. *Nature Reviews Endocrinology*, 2020, **16**(1): 45-57.
- [3] Fu Q G, Malchi T, Carter L J, *et al.* Pharmaceutical and personal care products: from wastewater treatment into agro-food systems [J]. *Environmental Science & Technology*, 2019, **53**(24): 14083-14090.
- [4] 樊静静, 王赛, 唐金鹏, 等. 广州市流溪河水体中 6 种内分泌干扰物时空分布特征与环境风险 [J]. *环境科学*, 2018, **39**(3): 1053-1064.
Fan J J, Wang S, Tang J P, *et al.* Spatio-temporal patterns and environmental risk of endocrine disrupting chemicals in the Liuxi River [J]. *Environmental Science*, 2018, **39**(3): 1053-1064.
- [5] Tan R J, Liu R X, Li B, *et al.* Typical endocrine disrupting compounds in rivers of northeast China: occurrence, partitioning, and risk assessment [J]. *Archives of Environmental Contamination and Toxicology*, 2018, **75**(2): 213-223.
- [6] Wang Y W, Li Y, Hu A Y, *et al.* Monitoring, mass balance and fate of pharmaceuticals and personal care products in seven wastewater treatment plants in Xiamen City, China [J]. *Journal of Hazardous Materials*, 2018, **354**: 81-90.
- [7] Boxall A B A, Rudd M A, Brooks B W, *et al.* Pharmaceuticals and personal care products in the environment: what are the big questions? [J]. *Environmental Health Perspectives*, 2012, **120**(9): 1221-1229.
- [8] Bu Q W, Cao Y B, Yu G, *et al.* Identifying targets of potential concern by a screening level ecological risk assessment of human use pharmaceuticals in China [J]. *Chemosphere*, 2020, **246**: 125818.
- [9] Luo Z F, Tu Y, Li H P, *et al.* Endocrine-disrupting compounds in the Xiangjiang River of China: spatio-temporal distribution, source apportionment, and risk assessment [J]. *Ecotoxicology and Environmental Safety*, 2019, **167**: 476-484.
- [10] Wang W F, Ndungu A W, Wang J. Monitoring of endocrine-disrupting compounds in surface water and sediments of the Three Gorges reservoir region, China [J]. *Archives of Environmental Contamination and Toxicology*, 2016, **71**(4): 509-517.
- [11] Batt A L, Kincaid T M, Kostich M S, *et al.* Evaluating the extent of pharmaceuticals in surface waters of the United States using a National-scale Rivers and Streams Assessment survey [J]. *Environmental Toxicology and Chemistry*, 2016, **35**(4): 874-881.
- [12] Guruge K S, Goswami P, Tanoue R, *et al.* First nationwide investigation and environmental risk assessment of 72 pharmaceuticals and personal care products from Sri Lankan surface waterways [J]. *Science of The Total Environment*, 2019, **690**: 683-695.
- [13] Archer E, Petrie B, Kasprzyk-Hordern B, *et al.* The fate of pharmaceuticals and personal care products (PPCPs), endocrine disrupting contaminants (EDCs), metabolites and illicit drugs in

- a WWTW and environmental waters [J]. *Chemosphere*, 2017, **174**: 437-446.
- [14] Zhang M, Shi Y J, Lu Y L, *et al.* The relative risk and its distribution of endocrine disrupting chemicals, pharmaceuticals and personal care products to freshwater organisms in the Bohai Rim, China [J]. *Science of the Total Environment*, 2017, **590-591**: 633-642.
- [15] Zheng Q, Zhang R J, Wang Y H, *et al.* Occurrence and distribution of antibiotics in the Beibu Gulf, China: impacts of river discharge and aquaculture activities [J]. *Marine Environmental Research*, 2012, **78**: 26-33.
- [16] 马永存. 长江口低氧区及邻近海域浮游植物研究 [D]. 上海: 上海海洋大学, 2013.
Ma Y C. Study on phytoplankton of Changjiang estuary hypoxia area and the adjacent east China sea [D]. Shanghai: Shanghai Ocean University, 2013.
- [17] 张莉. 长江流域上中游典型区水土保持生态恢复分区及恢复效果研究 [D]. 重庆: 西南农业大学, 2005.
Zhang L. Study on effect of ecological restoration in soil and water conservation and division of exemplary area in middle and upper reaches of Changjiang River [D]. Chongqing: Southwest Agricultural University, 2005.
- [18] Zhang Y Q, Johnson A C, Su C, *et al.* Which persistent organic pollutants in the rivers of the Bohai Region of China represent the greatest risk to the local ecosystem? [J]. *Chemosphere*, 2017, **178**: 11-18.
- [19] Su C, Lu Y L, Johnson A C, *et al.* Which metal represents the greatest risk to freshwater ecosystem in Bohai Region of China? [J]. *Ecosystem Health and Sustainability*, 2017, **3**(2): e01260.
- [20] Donnachie R L, Johnson A C, Moeckel C, *et al.* Using risk-ranking of metals to identify which poses the greatest threat to freshwater organisms in the UK [J]. *Environmental Pollution*, 2014, **194**: 17-23.
- [21] Donnachie R L, Johnson A C, Sumpter J P. A rational approach to selecting and ranking some pharmaceuticals of concern for the aquatic environment and their relative importance compared with other chemicals [J]. *Environmental Toxicology and Chemistry*, 2016, **35**(4): 1021-1027.
- [22] Johnson A C, Donnachie R L, Sumpter J P, *et al.* An alternative approach to risk rank chemicals on the threat they pose to the aquatic environment [J]. *Science of The Total Environment*, 2017, **599-600**: 1372-1381.
- [23] Johnson A C, Jürgens M D, Su C, *et al.* Which commonly monitored chemical contaminant in the Bohai region and the Yangtze and Pearl Rivers of China poses the greatest threat to aquatic wildlife? [J] *Environmental Toxicology and Chemistry*, 2018, **37**(4): 1115-1121.
- [24] Li Y, Zhang L Y, Liu X S, *et al.* Ranking and prioritizing pharmaceuticals in the aquatic environment of China [J]. *Science of The Total Environment*, 2019, **658**: 333-342.
- [25] Thorpe K L, Cummings R I, Hutchinson T H, *et al.* Relative potencies and combination effects of steroidal estrogens in fish [J]. *Environmental Science & Technology*, 2003, **37**(6): 1142-1149.
- [26] Holbech H, Kinnberg K, Petersen G I, *et al.* Detection of endocrine disruptors: evaluation of a fish sexual development test (FSDT) [J]. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 2006, **144**(1): 57-66.
- [27] Wang S, Zhu Z L, He J F, *et al.* Steroidal and phenolic endocrine disrupting chemicals (EDCs) in surface water of Bahe River, China; distribution, bioaccumulation, risk assessment and estrogenic effect on *Hemiculter leucisculus* [J]. *Environmental Pollution*, 2018, **243**: 103-114.
- [28] 陈晓雯, 赵建亮, 刘有胜, 等. 长江中下游环境激素效应的污染特征及生态风险 [J]. *生态毒理学报*, 2016, **11**(3): 191-203.
Chen X W, Zhao J L, Liu Y S, *et al.* Occurrence and ecological risks of hormonal activities in the middle and lower reaches of Yangtze River [J]. *Asian Journal of Ecotoxicology*, 2016, **11**(3): 191-203.
- [29] Brain R A, Johnson D J, Richards S M, *et al.* Effects of 25 pharmaceutical compounds to *Lemna gibba* using a seven-day static-renewal test [J]. *Environmental Toxicology and Chemistry*, 2004, **23**(2): 371-382.
- [30] Xu W H, Zhang G, Zou S C, *et al.* A preliminary investigation on the occurrence and distribution of antibiotics in the Yellow River and its tributaries, China [J]. *Water Environment Research*, 2009, **81**(3): 248-254.
- [31] Zhang Q Q, Ying G G, Pan C G, *et al.* Comprehensive evaluation of antibiotics emission and fate in the river basins of China: source analysis, multimedia modeling, and linkage to bacterial resistance [J]. *Environmental Science & Technology*, 2015, **49**(11): 6772-6782.
- [32] Yang Y Y, Cao X H, Zhang M M, *et al.* Occurrence and distribution of endocrine-disrupting compounds in the Honghu Lake and East Dongting Lake along the Central Yangtze River, China [J]. *Environmental Science and Pollution Research*, 2015, **22**(22): 17644-17652.
- [33] 陈政宏, 郭敏, 徐怀洲, 等. 太湖表层水体及沉积物中双酚 A 类似物的分布特征及潜在风险 [J]. *环境科学*, 2017, **38**(7): 2793-2800.
Chen M H, Guo M, Xu H Z, *et al.* Distribution characteristics and potential risk of bisphenol analogues in surface water and sediments of Lake Taihu [J]. *Environmental Science*, 2017, **38**(7): 2793-2800.
- [34] 朴海涛. 京杭运河及沿岸区域地表水中药物及个人护理品污染地理分布特征及来源辨析 [D]. 北京: 中国地质科学院, 2017.
Piao H T. Distribution and source analysis of pharmaceutical and personal care products in the Grand Canal and related areas [D]. Beijing: Chinese Academy of Geological Sciences, 2017.
- [35] 高月, 李杰, 许楠, 等. 汉江水相和沉积物中药品和个人护理品 (PPCPs) 的污染水平与生态风险 [J]. *环境化学*, 2018, **37**(6): 1706-1719.
Gao Y, Li J, Xu N, *et al.* Pollution levels and ecological risks of PPCPs in water and sediment samples of Hanjiang River [J]. *Environmental Chemistry*, 2018, **37**(6): 1706-1719.
- [36] Ma X Q, Wan Y J, Wu M Y, *et al.* Occurrence of benzophenones, parabens and triclosan in the Yangtze River of China, and the implications for human exposure [J]. *Chemosphere*, 2018, **213**: 517-525.
- [37] Liu Y H, Zhang S H, Song N H, *et al.* Occurrence, distribution and sources of bisphenol analogues in a shallow Chinese freshwater lake (Taihu Lake): implications for ecological and human health risk [J]. *Science of the Total Environment*, 2017, **599-600**: 1090-1098.
- [38] Wu M H, Xie D G, Xu G, *et al.* Benzophenone-type UV filters in surface waters: an assessment of profiles and ecological risks in Shanghai, China [J]. *Ecotoxicology and Environmental Safety*, 2017, **141**: 235-241.
- [39] Zhou L J, Wu Q L, Zhang B B, *et al.* Occurrence,

- spatiotemporal distribution, mass balance and ecological risks of antibiotics in subtropical shallow Lake Taihu, China [J]. *Environmental Science: Processes & Impacts*, 2016, **18**(4): 500-513.
- [40] Liu Y H, Zhang S H, Ji G X, *et al.* Occurrence, distribution and risk assessment of suspected endocrine-disrupting chemicals in surface water and suspended particulate matter of Yangtze River (Nanjing section) [J]. *Ecotoxicology and Environmental Safety*, 2017, **135**: 90-97.
- [41] Yao L L, Wang Y X, Tong L, *et al.* Occurrence and risk assessment of antibiotics in surface water and groundwater from different depths of aquifers: A case study at Jiangnan Plain, central China [J]. *Ecotoxicology and Environmental Safety*, 2017, **135**: 236-242.
- [42] Liu D, Liu J N, Guo M, *et al.* Occurrence, distribution, and risk assessment of alkylphenols, bisphenol A, and tetrabromobisphenol A in surface water, suspended particulate matter, and sediment in Taihu Lake and its tributaries [J]. *Marine Pollution Bulletin*, 2016, **112**(1-2): 142-150.
- [43] 丁惠君. 鄱阳湖水环境抗生素污染特征及典型抗生素的吸附和降解研究[D]. 武汉: 武汉大学, 2018.
Ding H J. Study on the characteristics of antibiotics in Poyang Lake and the adsorption and degradation of typical antibiotics [D]. Wuhan: Wuhan University, 2018.
- [44] Ma R X, Wang B, Lu S Y, *et al.* Characterization of pharmaceutically active compounds in Dongting Lake, China: occurrence, chiral profiling and environmental risk [J]. *Science of the Total Environment*, 2016, **557-558**: 268-275.
- [45] Wang Z, Du Y, Yang C, *et al.* Occurrence and ecological hazard assessment of selected antibiotics in the surface waters in and around Lake Honghu, China [J]. *Science of The Total Environment*, 2017, **609**: 1423-1432.
- [46] Peng Y, Fang W D, Krauss M, *et al.* Screening hundreds of emerging organic pollutants (EOPs) in surface water from the Yangtze River Delta (YRD): occurrence, distribution, ecological risk [J]. *Environmental Pollution*, 2018, **241**: 484-493.
- [47] Lin H J, Chen L L, Li H P, *et al.* Pharmaceutically active compounds in the Xiangjiang River, China: Distribution pattern, source apportionment, and risk assessment [J]. *Science of the Total Environment*, 2018, **636**: 975-984.
- [48] Xie Z X, Lu G H, Yan Z H, *et al.* Bioaccumulation and trophic transfer of pharmaceuticals in food webs from a large freshwater lake [J]. *Environmental Pollution*, 2017, **222**: 356-366.
- [49] Lu J, Li H P, Luo Z F, *et al.* Occurrence, distribution, and environmental risk of four categories of personal care products in the Xiangjiang River, China [J]. *Environmental Science and Pollution Research*, 2018, **25**(27): 27524-27534.
- [50] Yan M T, Xu C, Huang Y M, *et al.* Tetracyclines, sulfonamides and quinolones and their corresponding resistance genes in the Three Gorges Reservoir, China [J]. *Science of the Total Environment*, 2018, **631-632**: 840-848.
- [51] 封丽, 程艳茹, 封雷, 等. 三峡库区主要水域典型抗生素分布及生态风险评估 [J]. *环境科学研究*, 2017, **30**(7): 1031-1040.
Feng L, Cheng Y R, Feng L, *et al.* Distribution of typical antibiotics and ecological risk assessment in main waters of Three Gorges Reservoir area [J]. *Research of Environmental Sciences*, 2017, **30**(7): 1031-1040.
- [52] Wang Y Q, Liu Y, Lu S Y, *et al.* Occurrence and ecological risk of pharmaceutical and personal care products in surface water of the Dongting Lake, China-during rainstorm period [J]. *Environmental Science and Pollution Research*, 2019, **26**(28): 28796-28807.

CONTENTS

Industrial Volatile Organic Compounds (VOCs) Emission Inventory in China	LIANG Xiao-ming, SUN Xi-bo, XU Jian-tie, <i>et al.</i> (4767)
Contribution of Emissions from Cement to Air Quality in China	TANG Ling, XUE Xiao-da, BO Xin, <i>et al.</i> (4776)
Audit Indicators and Suggested Ranges for Data Validation of Chemical Components in Ambient PM _{2.5} : A Case Study of the Yangtze River Delta	 ZHANG Zhe, QIAO Li-ping, ZHOU Min, <i>et al.</i> (4786)
Size Distribution of Particulate Chemical Components in Nanjing Jiangbei New Area	ZHANG Yu-xiu, YU Xing-na, LIU Si-jia, <i>et al.</i> (4803)
Seasonal Chemical Composition Characteristics and Source Apportionment of PM _{2.5} in Zhengzhou	ZHANG Jian-fei, JIANG Nan, DUAN Shi-guang, <i>et al.</i> (4813)
Characteristics, Sources, and Health Risks of Elements in PM _{2.5} in Shanxi University Town	LI Xue-mei, MU Ling, TIAN Mei, <i>et al.</i> (4825)
Spatio-temporal Evolution of PM _{2.5} Concentration During 2000-2019 in China	XIA Xiao-sheng, WANG Jun-hong, SONG Wei-dong, <i>et al.</i> (4832)
Differences in Pollution Characteristics Under the Southerly and Easterly Wind in Beijing	YIN Xiao-mei, QIAO Lin, ZHU Xiao-wan, <i>et al.</i> (4844)
Transport Characteristics of Air Pollution in Tianjin Based on Weather Background	CAI Zi-ying, YAN Xu, HAN Su-qin, <i>et al.</i> (4855)
Influence of Meteorological Conditions on Ozone Pollution at Shangdianzi Station Based on Weather Classification	WU Jin, LI Chen, MA Zhi-qiang, <i>et al.</i> (4864)
Accurate Identification of Ambient VOCs Emission Sources in an Industrial Park Using On-Line Monitoring Data	SUN Jia-ning, XIE Shao-dong (4874)
Characteristics, Atmospheric Reactivity, and Source Apportionment of Ambient Volatile Organic Compounds in Wuhu	GAO Kang, ZHANG Hui, LIU Meng-di, <i>et al.</i> (4885)
Water Quality Response to Landscape Pattern at Different Spatial Scales in Baihe River in the Upper Reaches of the Miyun Reservoir	ZHANG Wei-wei, LI Xiao-na, WANG Chao, <i>et al.</i> (4895)
Concentration and Carbon Isotope Composition of DOC and DIC in the Miyun Reservoir Watershed in Summer	CHEN Ji-ji, GUO Jing, XU Su-shi, <i>et al.</i> (4905)
Variation and Driving Factors of Black Water Event Intensity in Lake Taihu During 2009 to 2018	XIN Hua-rong, ZHU Guang-wei, WANG Xue-song, <i>et al.</i> (4914)
Hydrogeochemical Investigations of Groundwater in the Lingbei Area, Leizhou Peninsula	ZHANG Hong-xin, WU Ya, LUO Wei-yu, <i>et al.</i> (4924)
Source of Groundwater Nitrate in Luanping Basin Based on Multi-environment Media Nitrogen Cycle and Isotopes	SUN Hou-yun, WEI Xiao-feng, JIA Feng-chao, <i>et al.</i> (4936)
Metal Distributions and Human Health Risk Assessments on Waters in the Huixian Karst Wetland, China	LI Jun, ZOU Sheng-zhang, LIANG Yong-ping, <i>et al.</i> (4948)
Absorption and Three Dimensional Fluorescence Spectra of CDOM in the Water of the Taibaishan Nature Reserve	ZHANG Wen-hao, ZHAO Duo-lin, WANG Xiao-yu, <i>et al.</i> (4958)
Effect of Heavy Rainfall on Nitrogen and Phosphorus Concentrations in Rivers at River-net Plain	LIAN Xin-qiao, ZHU Guang-wei, YANG Wen-bin, <i>et al.</i> (4970)
Risk Ranking of Endocrine Disrupting Compounds, Pharmaceuticals, and Personal Care Products in the Aquatic Environment of the Yangtze River Basin	SU Chao, CUI Yan (4981)
Sources and Ecological Risk Assessment of Polycyclic Aromatic Hydrocarbons in the Jiaxing River Network	YANG Yan-mei, ZHAO Hang-chen, MENG Rui, <i>et al.</i> (4989)
Impact of a Sewage Treatment Plant on the Accumulation of Microplastics in Freshwater Organisms in the Lijiang River of the Guilin Urban Section	 XIE Yuan-shan, ZHANG Qing-ke, ZHANG Li-shan, <i>et al.</i> (4999)
Distribution of Antibiotic Concentration in Domestic Wastewater Treatment Facilities in Villages and Towns	LU Ya-nan, GUO Ya-ni, WANG Kun, <i>et al.</i> (5008)
Microbial Diversity and Influencing Factors in a Small Watershed in Winter	ZHU Jin-shan, QIN Hai-lan, SUN Qi-yao, <i>et al.</i> (5016)
Characteristics of Organic Matter in Sediments During the Thermal Stratification of a Reservoir and Effects on an Aerobic Denitrifier	WANG Tong, WEN Gang, HUANG Ting-lin, <i>et al.</i> (5027)
Microbial Diversity Analysis of WWTPs Based on Hybrid-MBBR Process in a Low Temperature Season in the Yangtze River Delta	HAN Wen-jie, WU Di, ZHOU Jia-zhong, <i>et al.</i> (5037)
Effects of Phytoplankton Classifications on the Relationship Between Phytoplankton Community and Environment in Summer Subtropical Reservoirs, Southern China	 GUO Fang, GU Ji-Guang, ZHAO Jian, <i>et al.</i> (5050)
Remote Sensing Classification of Urban Black-odor Water Based on Decision Tree	LI Ling-ling, LI Yun-mei, LÜ Heng, <i>et al.</i> (5060)
Enhanced Membrane Anti-fouling Ability of Anaerobic Membrane Bioreactor Equipped with Bio-electrochemical System Using Nano-zero-valent Iron and Its Regulation Mechanism	 PAN Yang, NIU Cheng-xin, ZHI Zhong-xiang, <i>et al.</i> (5073)
Effects of Temperature and Substrate Concentration on N ₂ O Release of ANAMMOX Process	WANG Si-ke, YU Heng, ZUO Jian-e (5082)
Characteristics of Nutrient Removal in a Pilot-scale A ² /O with Mixture of Sludge Fermentation Liquor and Tail Water as External Carbon Source	 RONG Yi, SHAO Jin-bo, YU You-cheng, <i>et al.</i> (5089)
Effect of Different Ratios of Anaerobic Time and Aeration Time on the Operation of a Continuous-Flow Reactor with Partial Nitrification Granules	 ZHANG Jie, LAO Hui-mei, LI Dong, <i>et al.</i> (5097)
Effects of Mesophilic Anaerobic Digestion and Thermophilic Anaerobic Digestion on the Risk and Stability of Heavy Metals in Sludge	TIAN Zhong-ke, WANG Fen, YAN Zhao (5106)
Quantitative Evaluation of Human Health Risk of Heavy Metals in Soils Based on Positive Matrix Factorization Model and Geo-statistics	HAN Lin, XU Xi-bo (5114)
Distribution and Ecological Risk of Heavy Metals in the Soil of Redevelopment Industrial Sites	SHEN Cheng, LIU Fu-wen, WU Jian, <i>et al.</i> (5125)
Investigating Adsorption of Naphthalene to Organo-Modified Montmorillonites and Influencing Factors	HOU Jing-wen, SHI Wei-lin, GAO Juan (5133)
Effects of Ferrous Sulfate and Ferric Nitrate on Cadmium Transportation in the Rhizosphere Soil-Rice System	LI Yi-chun, CHEN Yong, TANG Ming-deng, <i>et al.</i> (5143)
Effects of Intercropping of <i>Brassica chinensis</i> L. and <i>Tagetes patula</i> L. on the Growth and Cadmium Accumulation of Plants	YAN Xiu-xiu, XU Ying-ming, WANG Lin, <i>et al.</i> (5151)
Effects of Heavy Metal-immobilizing Bacteria on Reducing Wheat Uptake Cd by Regulation Soil Iron Oxides	WANG Xiao-yu, HAN Hui (5160)
Sustained Effects of Remediation Materials on Soil Copper Remediation Under Oil-Rice Rotation	ZHAO Ying-ying, LI Hong-chuan, HU Hong-xiang, <i>et al.</i> (5168)
Impacts of Nitrogen Application on Ammonia Volatilization During Maize Season in Northern China	ZHANG Wei, NI Bang, XU Xiu-chun, <i>et al.</i> (5176)
Effect of Biochar on Changes of the Temperature Sensitivity of Soil Respiration and Bacterial Community Structure	CAO Kun-kun, ZHANG Sha-sha, HU Xue-yu, <i>et al.</i> (5185)
Effects of Heavy Metal Contents on Phyllosphere and Rhizosphere Fungal Communities for <i>Bothriochloa ischaemum</i> in Copper Tailings Area	 JIA Tong, GUO Ting-yan, WANG Rui-hong, <i>et al.</i> (5193)
Advances in Microbe-Derived Taste and Odor Compounds in Typical Aquatic Ecosystems	ZHANG Hai-han, MIAO Yu-tian, HUANG Ting-lin, <i>et al.</i> (5201)